

Central Mandibular Nerve Sheath Myxoma

Narges Ghazi¹, Jahanshah Salehinejad², Shapoor Yaripour³, Arash Dehghan⁴,
Elham Baradaran Nasser⁵, Mehdi Shahabinejad⁶

¹Oral & Maxillofacial Diseases Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

²Department of Oral and Maxillofacial Pathology, Mashhad Dental School, Mashhad, Iran

³Department of Oral and Maxillofacial Surgery, Hamedan Dental School Hamedan, Iran

⁴Department of General and Clinical Pathology, Hamedan University of Medical Science, Hamedan, Iran

⁵Department of Medicine, Mashhad Dental School, Mashhad, Iran

⁶Department of Oral and Maxillofacial Pathology, Mashhad Dental School, Mashhad, Iran

Received 16 October 2017 and Accepted 11 December 2017

Abstract

Nerve sheath myxoma has been described as a rare neural tumor arising from Schwann cells. It is observed most frequently in the central area of the face, neck and upper extremities. In the past the term neurothekeoma was used as synonym for nerve sheath myxoma but according to new reports, they are separate entities which can be confirmed by immunohistochemistry as in our case. Oral involvement of this tumor is extremely rare. Here, we present an unusual case of nerve sheath myxoma in the mandible of a 22-year old female patient. This case appears to be the first myxomatous variant which is centrally located in the mandible.

Keywords: nerve sheath myxoma, schwann cells, neurothekeoma, mandible.

Introduction

Nerve sheath myxoma (NSM) is an uncommon, benign nerve sheath tumor which was described in 1969 as a distinct pathologic entity (1-2). In the past the term neurothekeoma was used as synonym for nerve sheath myxoma but according to new reports they are separate entities which can be confirmed by immunohistochemistry (3). The microscopic differential diagnosis of nerve sheath myxoma include myxoid neurofibroma and myxoid neurothekeoma.

neurothekeoma is S100 negative while nerve sheath myxoma is S100 positive as in our case.

It is observed most frequently in the central area of the face, neck and upper extremities. A slight female predilection was reported (1.8:1) with the mean age of 25 years (4).

Histologically, it is characterized by the presence of epithelioid and spindle cells arranged in well-formed micronodules. Myxoid, mixed and cellular are the three histologic variants of NSM (4). The treatment of choice is wide local excision with free surgical margins and it has local recurrence rate of 50% but no metastasis or malignant transformations has been reported.

Oral involvement is extremely rare. Here, we present an unusual case of nerve sheath myxoma centrally located in the mandible of a young patient.

Ghazi N, Salehinejad J, Yaripour S, Dehghan D, Baradaran Nasser E, Shahabinejad M. Central mandibular nerve sheath myxoma. *J Dent Mater Tech* 2018; 7(1): 49-52.

Case report

A 22-year old female presented with a chief complaint of left mandibular swelling in the area of tooth #38 that had been started two weeks ago. The patient also had numbness of left side of lower lip and pain during chewing.

Clinical examination revealed a firm swelling, present on the left mandibular angle which was extended intraorally in the area of partially erupted tooth #38 (Fig. 1).




Figure 1. Gross appearance of resected mandible

Panoramic radiograph demonstrated a well- defined unilocular radiolucency extending from distal of tooth #37 up to the inferior border of condyle and coronoid causing obliteration of mandibular canal and thinning of inferior border of mandible (Fig. 2).

Computed tomography (CT) and 3D images showed a 44×22mm lesion with massive bone destruction and thinning of mandibular border which extends medially. A provisional diagnosis of KOT (keratocystic odontogenic tumor), ameloblastoma and myxoma was made (Fig. 2).




Figure 2. Panoramic radiograph and 3D images demonstrated a well- defined unilocular radiolucency with massive bone destruction and thinning of mandibular border

After incisional biopsy, the specimen submitted to the Pathology Department for histologic examination. Histopathological examination revealed myxomatous tissue consist of fine and scarce collagen fibers and few

fibroblast together with sections of blood vessels. According to histopathological findings an initial diagnosis of myxoma was made (Fig 3).




Figure 3. Photomicrograph showing A, B: Round to oval cells along with spindle cells with wavy nuclei against a myxoid background, H&E, original magnification, A: 100x, B:400x, C: Mast cell in 100x and, D: 400x

Regarding the extension of the lesion and histopathologic evaluation hemimandibulectomy was performed. The gross examination demonstrated a section of mandible measuring 8.5 ×4×2.5cm with irregular surface and boldging in some areas. (Fig. 1)

Microscopic examination showed myxomatous tissue consist of fine and scarce collagen fibers together with proliferation of small, spindle shape cells with wavy nuclei similar to Schwann cells .

Immunohistochemical examination of markers including vimentin, S-100, and neurofilament confirmed the diagnosis of nerve sheath myxoma (Fig. 4).

Intense positivity for vimentin, scattered positivity for S-100 and very scant positivity of neurofilament specially in the area of nerve bundles was present. (Fig. 4). Patient follow up had been done a year after and she was free of any recurrences.




Figure 4: Photomicrograph showing A,B: S-100 imunoreactivity A:400x, B:100x, C: Vimentine immunoreactivity, 100x.

Discussion

Nerve sheath myxoma is a rare neural tumor arising from Schwann cells. Rodriguez and el-Naggar reported that the mean age of patients with oral nerve sheath myxoma is 33 years (5).

Oral involvement of this tumor is extremely rare. Tongue is the most common site which has been

reported by several authors (6-8). Involvement of retromolar area (9), palate (10-11) and buccal mucosa (1, 5, 9) have been also reported. Intra-osseous nerve sheath myxoma of the jaw is very rare. kaltman et al. reported a cellular varient of mandibular NSM and it was S-100 negative (12).

Our case is myxomatous varient and is the first case which is centrally located in mandible. In our case

the neoplastic elements demonstrated S-100 immunoreactivity which according to Argenyi et al. the myxomatous variant is S-100 positive (13). Positive reactivity for S-100 has been also reported by several authors (3,8,11).

Histopathological differential diagnosis for NSM are myxoid neurothekeoma, neurilemmoma, myxoid neurofibroma and oral myxoma (14). In myxoid neurothekeoma the myxoid area is located at the periphery of tumor but in our case it is more of central location and myxoid neurothekeoma is S100 negative.(3) Both of NSM and neurilemmoma arise from the nerve sheath and may show positive reactivity for S100. Myxoid appearance is rarely seen in neurilemmoma.

Although areas of increased cellularity may be seen in NSM similar to neurilemmoma, Verocay body formation is not observed in NSM (14).

The histopathologic feature of myxoid neurofibroma is a circumscribed and unencapsulated tumor with slender, closely aggregated fascicles of Schwann cells. The myxoid matrix is confined within fascicles in NSM, whereas in neurofibroma it usually involves both fascicles and surrounding tissue (8). It should be noted that lobular pattern and S100 immunoreactivity are not seen in oral myxomas (5).

The treatment of choice for NSM is local excision (8). Regarding the extension of the lesion and histopathologic evaluation hemimandibulectomy was performed for the present case.

References

1. Barrett AW, Suhr M. Cellular neurothekeoma of the oral mucosa. *Oral Oncol* 2001; 37(8): 660-4.
2. Fetsch JF, Laskin WB, Hallman JR, Lupton GP, Miettinen M. Neurothekeoma: an analysis of 178 tumors with detailed immunohistochemical data and long-term patient follow-up information. *Am J Surg Pathol* 2007; 31(7): 1103-14.
3. Amoolya Bhat, Aparna Narasimha, Vijaya C, Sundeep VK. Nerve Sheath Myxoma: Report of a Rare Case. *J Clin Diagn Res* 2015 Apr; 9(4): ED07–ED09.
4. Beena VT, Deepthi PV, Padmakumar SK, Sivakumar R. Neurothekeoma of oral cavity: a rare

- case report. *Int J Scientific Stud* 2015; 12(2): 208-10.
5. Rodriguez-Peralto JL, el-Naggar AK. Neurothekeoma of the oral cavity: case report and review of the literature. *J Oral Maxillofac Surg* 1992; 50(11): 1224-6.
6. Mincer HH, Spears KD. Nerve sheath myxoma in the tongue. *Oral Surg Oral Med Oral Pathol* 1974; 37(3) :428-30.
7. Yamamoto H, Kawana T. Oral nerve sheath myxoma. Report of a case with findings of ultrastructural and immunohistochemical studies. *Acta Pathol Jpn* 1988; 38(1): 121-7.
8. Peñarrocha M, Bonet J, Minguez JM, Vera F. Nerve sheath myxoma (neurothekeoma) in the tongue of a newborn. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2000; 90(1): 74-7.
9. Sist TC Jr, Greene GW Jr. Benign nerve sheath myxoma: light and electron microscopic features of two cases. *Oral Surg Oral Med Oral Pathol* 1979; 47(5): 441-4.
10. Wright BA, Jackson D. Neural tumors of the oral cavity. A review of the spectrum of benign and malignant oral tumors of the oral cavity and jaws. *Oral Surg Oral Med Oral Pathol*. 1980 Jun; 49(6): 509-22.
11. Smith BC, Ellis GL, Meis-Kindblom JM, Williams SB. Ectomesenchymal chondromyxoid tumor of the anterior tongue. Nineteen cases of a new clinicopathologic entity. *Am J Surg Pathol* 1995; 19(5): 519-30.
12. Kaltman J, Best S, Kaltman S, Lopez E, Velez I. Intraosseous neurothekeoma of the mandible: A rare occurrence. *International journal of oral & maxillofacial pathology*; 2012; 3(3): 13-17.
13. Argenyi ZB, LeBoit PE, Santa Cruz D, Swanson PE, Kutzner H. Nerve sheath myxoma (neurothekeoma) of the skin: light microscopic and immunohistochemical reappraisal of the cellular variant. *J Cutan Pathol* 1993; 20: 294-303.
14. Chou LS, Hansen LS, Daniels TE. Christomas of the oral cavity: a review. *Oral Surg Oral Med Oral Pathol* 1991; 72(5): 584-93.

Corresponding Author:

Mehdi Shahabinejad,
Oral and maxillofacial pathology,
mashhad dental school, mashhad university of medical science, mashhad, iran
Tell: 05138829511
Email: shahabinejadm@mums.as.ir